arXiv:hep-ph/9909220v1 2 Sep 1999

UL–NTZ 18/99

Helicity amplitudes for the small–angle process $e^-e^\pm \to e^-\gamma\gamma + e^\pm$ with both photons along one direction and its cross channel

E.A. Kuraev [a,1], A. Schiller [b,2], V.G. Serbo [c,3],
B.G. Shaikhatdenov [a,4]

[a] *Joint Institute for Nuclear Research, Dubna, 141980, Russia*

[b] *Institut für Theoretische Physik and NTZ, Universität Leipzig, D-04109 Leipzig, Germany,*

[c] *Novosibirsk State University, Novosibirsk, 630090, Russia*

Abstract

We study the small–angle double bremsstrahlung in $e^-e^\pm$ scattering for a jet kinematics where both photons move along the electron direction. This region gives the main contribution to the cross section. We present analytic expressions for all 64 amplitudes with arbitrary helicity states of the initial and final leptons and the produced photons convenient for analytic and numerical studies. The accuracy of the obtained amplitudes is given omitting only terms of the order of m^2/E_j^2, θ_j^2 and $\theta_j m/E_j$. The helicity amplitudes for the cross channel $\gamma e^\pm \to \gamma e^+e^- + e^\pm$ are given. Several limits for the helicity amplitudes of hard or soft final particles are considered.

PACS: 12.20.-m; 13.85.Qk; 13.88.+e; 13.40.-f

Key words: Quantum electrodynamics, polarized leptons, polarized photons, jet kinematics

---

[1] E–mail: kuraev@thsun1.jinr.ru
[2] E–mail: schiller@tph204.physik.uni-leipzig.de
[3] E–mail: serbo@math.nsc.ru
[4] E–mail: sbg@thsun1.jinr.ru

Preprint submitted to Elsevier Preprint *2 September, 1999*

1 Introduction

Nowadays colliders and detectors are able to work with polarized particles. Therefore QED jet–like processes are required to be described in full detail including the helicities of all particles. Among other reactions, those inelastic processes whose cross sections do not fall with energy are of special interest. In particular, QED jets formed by two or three particles in reactions of the type

$$e^+e^- \to e^+\gamma + e^- \,, \quad e^+e^- \to e^+\gamma\gamma + e^- \,, \quad e^+e^- \to e^+e^+e^- + e^-$$

provide a realistic model for hadronic jets. The complete set of all QED jet–like tree processes up to order α^4 in the electromagnetic coupling have been listed in our previous paper [1]. For all of them except one we have obtained compact and simple analytic expressions for all helicity amplitudes to a high accuracy [2,3,1]. The process of double bremsstrahlung along one direction and its cross channel, discussed in the present paper, completes the calculation of all helicity amplitudes to order α^4.

Let us define by a jet kinematics in QED a high energy reaction in which the outgoing leptons and photons are produced within a small cone[5] relative to an axis given by their parental incoming lepton or photon. At high energies with the condition (p_1, p_2 are the incoming 4-momenta, m_j are the lepton masses)

$$s = 2p_1p_2 = 4E_1E_2 \gg m_j^2 \tag{1}$$

the dominant contribution to the non-decreasing cross sections is given by the region of scattering angles θ_j which is much smaller than unity though could be of the order of the typical emission angles m_j/E_j or larger:

$$m_j/E_j \lesssim \theta_j \ll 1 \ . \tag{2}$$

In this kinematic region all processes have the form of two–jet processes with an exchange of a single virtual photon γ^* in the t–channel (see Fig. 1).

In the kinematic region (1), (2) it is possible to obtain compact and relatively simple analytic expressions for all helicity amplitudes of the jet-like QED processes to a high accuracy omitting terms of the relative order of

---

[5] The opening angle of the cone is characterized by the maximal polar angles of the produced particles.



Fig. 1. Generic block diagram $ee \to$ two jets.

$$\frac{m_j^2}{E_j^2}\,, \quad \theta_j^2\,, \quad \frac{m_j}{E_j}\,\theta_j \tag{3}$$

only.

The amplitude M_{fi} corresponding to the diagram of Fig. 1 can be presented in the form

$$M_{fi} = M^{\mu}_{\rm up}\,\frac{g_{\mu\nu}}{k^2}\,M^{\nu}_{\rm down}\,, \tag{4}$$

where $M^{\mu}_{\rm up}$ and $M^{\nu}_{\rm down}$ are the amplitudes of the upper and lower block of Fig. 1, respectively, and $(-g_{\mu\nu}/2)$ is the density matrix of the virtual photon. The transition amplitude $M_{\rm up}$ describes the scattering of an incoming particle (in our case a lepton) with a virtual photon of "mass" squared k^2 and polarization 4–vector $e = k_\perp/\sqrt{-k_\perp^2}$ ($k_\perp$ is the 4–vector transverse to p_1 and p_2) to some QED final state in the jet kinematics of Eqs. (1), (2) (similar for $M_{\rm down}$).

To introduce some notations for e^-e^+ collisions we use the block diagram of Fig. 1 as an example. We work in a reference frame in which the initial particles with 4–momenta p_1 and p_2 perform a head–on collision with energies E_1 and E_2 of the same order. The z–axis is chosen along the momentum of the first initial particle, the azimuthal angles are denoted by φ_j (they are referred to one fixed x-axis). Exploiting the freedom in choosing a general phase factor, we put the initial azimuthal angle of the electron φ_1 and that of the positron φ_2 equal to zero:

$$\varphi_1 = \varphi_2 = 0\,. \tag{5}$$

It is convenient to introduce "the almost light–like" 4–vectors p and p'

$$\begin{aligned} &p = p_1 - \frac{m^2}{s}\,p_2\,, \quad p' = p_2 - \frac{m^2}{s}\,p_1\,, \quad p_1^2 = p_2^2 = m^2\,, \\ &p^2 = p'^2 = \frac{m^6}{s^2} \approx 0\,, \quad s = 2p_1p_2 = 2pp' + 3\frac{m^4}{s}\,. \end{aligned} \tag{6}$$

With accuracy (3) the matrix $g_{\mu\nu}$ from Eq. (4) can be transformed to the form

$$g_{\mu\nu} \to \frac{2}{s} p'_\mu p_\nu \,, \tag{7}$$

(for details, see [4], §4.8.4) which results in the factorized representation for the amplitude

$$M_{fi} = \frac{s}{k^2} J_{\rm up} J_{\rm down} \,. \tag{8}$$

The vertex factors $J_{\rm up,down}$ are given by the block amplitudes $M^\mu_{\rm up}$ and $M^\nu_{\rm down}$

$$J_{\rm up} = \frac{\sqrt{2}}{s} M^\mu_{\rm up}\, p'_\mu \,, \quad J_{\rm down} = \frac{\sqrt{2}}{s} M^\nu_{\rm down}\, p_\nu \,. \tag{9}$$

The quantities $J_{\rm up}$ and $J_{\rm down}$ can be calculated in the limit $s \to \infty$ assuming that the energy fractions and transverse momenta of the final particles are finite.

We study the double bremsstrahlung in one direction for small angle $e^- e^\pm$ scattering

$$e^-(p_1) + e^\pm(p_2) \to e^-(p_3) + \gamma(k_1) + \gamma(k_2) + e^\pm(p_4) \,. \tag{10}$$

For unpolarized particles this process has been discussed in Refs. [5] and [6] where the differential cross sections and photon spectra were calculated in the approximation of classical currents and equivalent photons. In Refs. [6,7] the matrix element squared for the unpolarized case has been found to the accuracy (3). The helicity amplitudes of the process (10) for large angle scattering have been calculated in Refs. [8,9]. The recent interest to multiphoton emission is related to the exact calculation of Bhabha scattering needed for LEP luminosity calibration (see, for example, Refs. [10]).

In Born approximation the process (10) (with photons not only along one direction) is described by 32 Feynman diagrams. In the considered kinematics (forward scattering at high energies) only 16 scattering–type diagrams are relevant since the contribution to the cross section which does not decrease with the energy arises from those Feynman diagrams with a virtual photon exchange in the t–channel. Restricting to the case in which both photons are emitted along the electron (p_1) direction, only six diagrams are relevant. Three of them are shown in Fig. 2.



Fig. 2. Feynman diagrams for double bremsstrahlung in e^-e^+ scattering, diagrams with $k_1 \leftrightarrow k_2$ have to be added.

2 The double bremsstrahlung helicity amplitudes in the jet kinematics

2.1 The Sudakov 4-vector decomposition and kinematic invariants

The amplitude for the double forward bremsstrahlung of the process (10) can be written with accuracy (1), (2) in the factorized form (for definiteness the Bhabha scattering is considered, see Fig. 2, where the momenta and helicities of all particles are indicated)

$$\begin{aligned} &M(e^-e^+ \to e^-\gamma\gamma + e^+) = \frac{s}{k^2} J_{\text{up}}(e^-\gamma^* \to e^-\gamma\gamma) J_{\text{down}}(e^+\gamma^* \to e^+)\,, \quad (11)\\ &k = p_4 - p_2 = p_1 - p_3 - k_1 - k_2\,. \end{aligned}$$

The vertex factor J_{down} has been calculated earlier [1]:

$$J_{\text{down}}(e^+_{\lambda_{\bar{e}2}}\gamma^* \to e^+_{\lambda_{\bar{e}4}}) = \sqrt{8\pi\alpha}\, \mathrm{e}^{-\mathrm{i}\,\lambda_{\bar{e}4}\,\varphi_4}\, \delta_{\lambda_{\bar{e}2},\lambda_{\bar{e}4}}\,, \qquad (12)$$

The quantity $\lambda_{\bar{e}2} = \pm 1/2$ ($\lambda_{\bar{e}4}$) denotes the helicities of the initial (final) positron, φ_4 is the azimuthal angle of the final e^+.

We have to calculate the upper vertex J_{up}. The almost light-like vectors p and p' have components $p = E_1(1,0,0,1)$, $p' = E_2(1,0,0,-1)$. It is convenient to use the Sudakov decomposition of any 4–vector along p and p' and transverse to them. For the momenta of the photons and the initial and final electron we have

$$\begin{aligned} &k_i = \alpha_i p' + x_i p + k_{i\perp}\,, \quad k_{i\perp}^2 = -\mathbf{k}_i^2\,, \quad (i = 1,2)\,,\\ &k = \alpha_k p' + \beta_k p + k_\perp\,, \quad k_\perp^2 = -\mathbf{k}^2\,, \qquad (13)\\ &p_1 = \frac{m^2}{s} p' + p\,, \quad p_3 = \alpha_3 p' + x_3 p + p_{3\perp}\,, \quad p_{3\perp}^2 = -\mathbf{p}_3^2\,, \end{aligned}$$

with the two–dimensional Euclidean vectors $\mathbf{k}$, $\mathbf{k}_i$ and $\mathbf{p}_3 = -\mathbf{k} - \mathbf{k}_1 - \mathbf{k}_2$

perpendicular to the beam axes, e.g.

$$\mathbf{k} = (k_x, k_y)\,. \tag{14}$$

In the kinematics of a jet moving close to the direction of the first electron the 2–vectors $\mathbf{k}_i$ and $\mathbf{p}_3$ have typical values of a few electron masses whereas $\mathbf{k}$ might be much smaller. The quantities $x_{1,2}$ and $x_3 = 1 - x_1 - x_2$ are the energy fractions of the emitted photons and the scattered electron

$$x_i = \frac{k_{i0}}{E_1}, \qquad x_3 = \frac{p_{30}}{E_1} = \frac{E_3}{E_1}\,. \tag{15}$$

They are supposed to be of the order of unity while α_i, α_3, α_k and β_k are small and vanish as $1/s$. Later on we often use complex circular components of a generic 2–vector transverse to the beam axes denoted by the corresponding Greek characters, e.g. for the 2–vector (14)

$$\kappa = k_x + \mathrm{i}\,k_y\,, \quad \kappa^* = k_x - \mathrm{i}\,k_y\,. \tag{16}$$

The on–shell conditions for the incoming electron and the outgoing electron and photons lead to

$$\begin{aligned} s\alpha_i &= \frac{\mathbf{k}_i^2}{x_i}\,, \quad s\alpha_3 = \frac{m^2 + \mathbf{p}_3^2}{x_3}\,, \\ s(\alpha_k + \alpha_1 + \alpha_2) &= -\frac{1}{x_3}\left[m^2(1 - x_3) + \mathbf{p}_3^2\right]\,. \end{aligned} \tag{17}$$

From the on–shell condition for the outgoing spectator (a positron in Bhabha scattering) $p_4^2 = (p_2 + k)^2 = m^2$ we find the expression

$$s\alpha_k\beta_k - \mathbf{k}^2 + s\beta_k + m^2\alpha_k = 0 \tag{18}$$

and present the momentum transfer squared (the 4–momentum of the virtual photon k) in the form

$$k^2 = s\alpha_k\beta_k - \mathbf{k}^2 = -\frac{1}{1 + \alpha_k}\left(\mathbf{k}^2 + m^2\alpha_k^2\right)\,. \tag{19}$$

Taking into account Eq. (19), we observe that the Sudakov parameter α_k is related to the invariant mass squared of the jet generated by the scattered electron and two accompanying photons

$$(p_3 + k_1 + k_2)^2 = (p_1 - k)^2 = m^2 - \mathbf{k}^2 - s\alpha_k\,. \tag{20}$$

From Eqs. (17), (18) and (19) explicit expressions for α_k and β_k can be derived.

After replacing all Sudakov parameters by the energy fractions x_i, x_3 and the 2-vectors $\mathbf{k}_i$ and $\mathbf{k}$, the kinematic invariants appearing in the upper vertex take the following form:

$$\begin{aligned}
a_i &\equiv -(p_1 - k_i)^2 + m^2 = \frac{1}{x_i}(m^2 x_i^2 + \mathbf{k}_i^2)\,,\\
b_i &\equiv (p_3 + k_i)^2 - m^2 = \frac{1}{x_i x_3}(m^2 x_i^2 + \mathbf{r}_i^2)\,,\\
a &\equiv -(p_1 - k_1 - k_2)^2 + m^2 = a_1 + a_2 - \frac{1}{x_1 x_2}\,(x_1\mathbf{k}_2 - x_2\mathbf{k}_1)^2\\
&= \frac{1}{x_1 x_2}\Big[x_1 x_2(1 - x_3)m^2 + x_2(1 - x_2)\mathbf{k}_1^2 + x_1(1 - x_1)\mathbf{k}_2^2 + 2x_1 x_2 \mathbf{k}_1\mathbf{k}_2\Big]\,,\\
b &\equiv (p_3 + k_1 + k_2)^2 - m^2 = b_1 + b_2 + \frac{1}{x_1 x_2}\,(x_1\mathbf{k}_2 - x_2\mathbf{k}_1)^2 \qquad (21)\\
&= \frac{1}{x_1 x_2 x_3}\Big[x_1 x_2(1 - x_3)m^2 + x_2(1 - x_2)(\mathbf{k}_1 + x_1\mathbf{k})^2\\
&\quad + x_1(1 - x_1)(\mathbf{k}_2 + x_2\mathbf{k})^2 + 2x_1 x_2(\mathbf{k}_1 + x_1\mathbf{k})(\mathbf{k}_2 + x_2\mathbf{k})\Big]\,,
\end{aligned}$$

where

$$\mathbf{r}_1 = x_1(\mathbf{k}_2 + \mathbf{k}) + (1 - x_2)\mathbf{k}_1\,, \quad \mathbf{r}_2 = x_2(\mathbf{k}_1 + \mathbf{k}) + (1 - x_1)\mathbf{k}_2 \tag{22}$$

or

$$\mathbf{r}_i = x_3\mathbf{k}_i - x_i\mathbf{p}_3\,. \tag{23}$$

Note that $b = -\mathbf{k}^2 - s\alpha_k$ and $b \to a/x_3$ in the limit $\mathbf{k} \to 0$.

Using the gauge invariance of an amplitude containing photons we can replace the polarization vector e_i of a photon i [6] with external momentum k_i by $e_i + \zeta_i k_i$. The arbitrary parameter ζ_i is chosen in such a way that the polarization vector does not contain a Sudakov projection onto the light–like vector p:

$$e_i \equiv e^{(\lambda_i)}(k_i) = \alpha_{e_i} p' + e_\perp^{(\lambda_i)}\,. \tag{24}$$

The transverse component $e_\perp$ which does not depend on the 4–momentum of the photon k_i is chosen as

---

[6] As usual, e_i (e_i^*) are used for incoming (outgoing) photons.

$$e_\perp^{(\lambda_i)} = -\frac{\lambda_i}{\sqrt{2}}(0, 1, \mathrm{i}\,\lambda_i, 0)\,, \quad e_\perp^{(\lambda_i)*} = -e_\perp^{(-\lambda_i)}\,. \tag{25}$$

The Sudakov parameters α_{e_i} are found from the conditions $e_i k_i = 0$:

$$s\alpha_{e_i} = \frac{2\mathbf{e}^{(\lambda_i)}\mathbf{k}_i}{x_i} = \frac{\sqrt{2}}{x_i}(-\delta_{\lambda_i,1}\kappa_i + \delta_{\lambda_i,-1}\kappa_i^*)\,. \tag{26}$$

Here κ_i and κ_i^* are the circular components of the vector $\mathbf{k}_i$ [compare Eq. (16)].

2.2 Helicity amplitudes for the upper vertex factor $J_{\mathrm{up}}(e^-\gamma^* \to e^-\gamma\gamma)$

Now all ingredients are prepared to calculate the helicity amplitudes for the upper vertex factor. We will use the following notation indicating the helicity states of the electron λ_{ei} and of the photons λ_i explicitly:

$$J_{\mathrm{up}} \equiv \frac{\sqrt{2}(4\pi\alpha)^{3/2}}{s}\,\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}}\,. \tag{27}$$

The matrix element $\mathcal{M}$ can be represented by the Feynman diagrams shown in Fig. 3. The amplitudes corresponding to that figure are



Fig. 3. Feynman diagrams for $e^-\gamma^* \to e^-\gamma\gamma$, diagrams with $k_1 \leftrightarrow k_2$ have to be added.

$$\mathcal{M} = (1 + \mathcal{P}_{12})Q\,, \qquad Q = \mathcal{M}_1 + \mathcal{M}_2 + \mathcal{M}_3\,, \tag{28}$$

where

$$\begin{aligned}
\mathcal{M}_1 &= \frac{1}{a_1 a}\bar{u}_3\hat{p}'(\hat{p}_1 - \hat{k}_1 - \hat{k}_2 + m)\hat{e}_2^*(\hat{p}_1 - \hat{k}_1 + m)\hat{e}_1^* u_1\,,\\
\mathcal{M}_2 &= -\frac{1}{a_1 b_2}\bar{u}_3\hat{e}_2^*(\hat{p}_1 - \hat{k}_1 - \hat{k} + m)\hat{p}'(\hat{p}_1 - \hat{k}_1 + m)\hat{e}_1^* u_1\,,\\
\mathcal{M}_3 &= \frac{1}{b_2 b}\bar{u}_3\hat{e}_2^*(\hat{p}_1 - \hat{k}_1 - \hat{k} + m)\hat{e}_1^*(\hat{p}_1 - \hat{k} + m)\hat{p}' u_1\,.
\end{aligned} \tag{29}$$

The permutation operator $\mathcal{P}_{12}$ for the photons is defined as

$$\mathcal{P}_{12} f(k_1, e_1; k_2, e_2) = f(k_2, e_2; k_1, e_1)\,, \qquad \mathcal{P}_{12}^2 = 1.$$

The quantity Q is gauge invariant with respect to the virtual photon k since all permutations of this photon are taken into account. Therefore Q is proportional to $k_\perp$ in the limit $k_\perp \to 0$. Indeed, using the relations

$$Q = p'_\mu Q^\mu\,, \quad k_\mu Q^\mu = (\alpha_k p' + \beta_k p + k_\perp)_\mu Q^\mu = 0\,, \tag{30}$$

we immediately obtain (neglecting the small contribution $\beta_k p_\mu Q^\mu \sim 1/s$)

$$Q = -\frac{k_{\perp\mu}}{\alpha_k} Q^\mu\,. \tag{31}$$

Now we transform the quantities $\mathcal{M}_j$ to such a form that in their sum Q this noticed low $k_\perp$ behaviour is present explicitly. The reason is that in that case all individual large (compared to $k_\perp$) contributions are cancelled. The first step is to use the Dirac equations $\hat{p}_1 u_1 = m u_1$, $\bar{u}_3 \hat{p}_3 = m \bar{u}_3$ and to rearrange the amplitudes $\mathcal{M}_j$ of Eq. (29) as follows:

$$\begin{aligned}
\mathcal{M}_1 &= \bar{u}_3 \left\{ \frac{s x_3}{a_1 a} \hat{e}_2^* (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* + \frac{1}{a_1 a} \hat{p}' \hat{k} \hat{e}_2^* (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* \right\} u_1\,, \\
\mathcal{M}_2 &= \bar{u}_3 \left\{ -\frac{s(1-x_1)}{a_1 b_2} \hat{e}_2^* (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* - \frac{1}{b_2} \hat{e}_2^* \hat{p}' \hat{e}_1^* \right. \\
&\qquad \left. + \frac{1}{a_1 b_2} \hat{e}_2^* \hat{k} \hat{p}' (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* \right\} u_1\,, \\
\mathcal{M}_3 &= \bar{u}_3 \left\{ \frac{s}{b_2 b} \hat{e}_2^* (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* - \frac{s}{b_2 b} \hat{e}_2^* \hat{k} \hat{e}_1^* \right. \\
&\qquad \left. - \frac{1}{b_2 b} \hat{e}_2^* (\hat{p}_3 + \hat{k}_2 + m) \hat{e}_1^* \hat{k} \hat{p}' \right\} u_1\,.
\end{aligned} \tag{32}$$

From these formulae we observe that the last terms in $\mathcal{M}_1, \mathcal{M}_2, \mathcal{M}_3$ are proportional to $k_\perp$ up to terms of the order of (3):

$$\hat{p}' \hat{k} = \hat{p}' (\alpha_k \hat{p}' + \beta_k \hat{p} + \hat{k}_\perp) = \hat{p}' \hat{k}_\perp = -\hat{k} \hat{p}'\,. \tag{33}$$

As a next step we note that the sum of the first three terms in Eqs. (32) is also proportional to $k_\perp$ since [see Eqs. (21)]

$$A \equiv \frac{x_3}{a_1 a} - \frac{1-x_1}{a_1 b_2} + \frac{1}{b_2 b}\,, \qquad A|_{k_\perp \to 0} = 0\,. \tag{34}$$

Finally we consider the sum of the second terms of the quantities $\mathcal{M}_2, \mathcal{M}_3$ given in Eqs. (32). Using the relations (21) and (20) one gets

$$-\frac{\hat{p}'}{b_2} - \frac{s(\alpha_k \hat{p}' + \hat{k}_\perp)}{b_2 b} = -\frac{s\hat{k}_\perp}{b_2 b} + \frac{\hat{p}'\mathbf{k}^2}{b_2 b}\,. \tag{35}$$

Therefore, from Eqs. (33), (34), (35) it is clearly seen that the discussed property (31)

$$(\mathcal{M}_1 + \mathcal{M}_2 + \mathcal{M}_3)\,|_{k_\perp \to 0} = 0$$

is obviously satisfied and the quantity $Q = \sum_{j=1}^{3} \mathcal{M}_j$ is a sum of terms explicitly proportional to $k_\perp$:

$$\begin{aligned} Q = \bar{u}_3 \Big\{ & As\,\hat{e}_2^*(\hat{p}_1 - \hat{k}_1 + m)\hat{e}_1^* + \frac{1}{a_1 a}\hat{p}'\hat{k}_\perp \hat{e}_2^*(\hat{p}_1 - \hat{k}_1 + m)\hat{e}_1^* \\ & - \frac{\mathbf{k}^2}{b_2 b}\hat{e}_2^*\hat{e}_1^*\hat{p}' - \frac{s}{b_2 b}\hat{e}_2^*\hat{k}_\perp \hat{e}_1^* \\ & + \frac{1}{a_1 b_2}\hat{e}_2^*\hat{k}_\perp \hat{p}'(\hat{p}_1 - \hat{k}_1 + m)\hat{e}_1^* - \frac{1}{b_2 b}\hat{e}_2^*(\hat{p}_3 + \hat{k}_2 + m)\hat{e}_1^*\hat{k}_\perp \hat{p}' \Big\} u_1\,. \end{aligned} \tag{36}$$

The subsequent calculations require to determine several bilinear spinor combinations which are listed in Appendix A.

As a result, we obtain the complete expression for the amplitude of double forward bremsstrahlung in a jet kinematics

$$M(e^- e^+ \to e^-\gamma\gamma + e^+) = \frac{32\pi^2\alpha^2}{k^2}\,\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}}\,\mathrm{e}^{-\mathrm{i}\,\lambda_{\bar{e}4}\,\varphi_4}\,\delta_{\lambda_{\bar{e}2},\lambda_{\bar{e}4}}\,. \tag{37}$$

All helicity amplitudes $\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}}$ are expressed via the complex circular components κ_i, κ and ρ_i of the vectors $\mathbf{k}_i$, $\mathbf{k}$ and $\mathbf{r}_i$ [see Eqs. (16) and (22)]

$$\kappa_i = k_{ix} + \mathrm{i}\,k_{iy}\,, \quad \kappa = k_x + \mathrm{i}\,k_y\,, \quad \rho_i = r_{ix} + \mathrm{i}\,r_{iy}\,, \quad i = 1, 2\,. \tag{38}$$

The spin non–flip amplitudes (with $\lambda_{e1} = \lambda_{e3} = +1/2 \equiv +$) being proportional to the factor

$$S_{\lambda_{e3}} = s\sqrt{x_3}\mathrm{e}^{\mathrm{i}\,\varphi_3 \lambda_{e3}}\,, \tag{39}$$

are equal to [7]

$$\begin{aligned}
\mathcal{M}^{++}_{++} &= 2S_+(1+\mathcal{P}_{12})\left\{A\frac{\kappa_1^*\rho_2^*}{x_1x_2x_3}-\frac{\kappa_1^*\kappa^*}{x_1a_1a}+\frac{\kappa^*\rho_2^*}{x_2x_3b_2b}\right\},\\
\mathcal{M}^{--}_{++} &= x_3S_+\left(\frac{\mathcal{M}^{++}_{++}}{S_+}\right)^*,\\
\mathcal{M}^{+-}_{++} &= -2S_+\left(C+\mathcal{P}_{12}D\right),\\
\mathcal{M}^{-+}_{++} &= \mathcal{P}_{12}\mathcal{M}^{+-}_{++},\\
C &= A\left(\frac{\kappa_1^*\kappa_2}{x_1x_2}-\frac{x_1}{x_3}m^2\right)-\frac{\kappa_1^*\kappa}{x_1a_1b_2}+\frac{x_1x_2\kappa^*\kappa+x_1\kappa^*\kappa_2+x_2\kappa_1^*\kappa}{x_1x_2b_2b},\\
D &= (1-x_1)\left(A\frac{\kappa_1\rho_2^*}{x_1x_2x_3}-\frac{\kappa_1\kappa^*}{x_1a_1a}+\frac{\kappa\rho_2^*}{x_2x_3b_2b}\right).
\end{aligned}\tag{40}$$

Another convenient form of C useful for the soft photon limit (see Section 3.1) and for the later crossing (Section 4) is

$$\begin{aligned}
C &= \frac{1-x_2}{1-x_1}D^*+\Delta,\\
\Delta &= -\frac{1}{a}+\frac{1}{b}+\frac{x_1x_2m^2+\kappa_1^*\rho_2}{x_3a_1b_2}+\frac{\kappa_1^*(x_2\kappa_1-x_1\kappa_2)}{x_1a_1a}-\frac{(x_2\kappa_1^*-x_1\kappa_2^*)\rho_2}{x_2x_3b_2b}.
\end{aligned}\tag{41}$$

Note that in the limit of a soft (second) photon the function $D\propto 1/x_2$ whereas Δ remains finite.

For the spin flip amplitudes (with $\lambda_{e1}=-\lambda_{e3}=+1/2$) proportional to mS_- we obtain

$$\begin{aligned}
\mathcal{M}^{++}_{+-} &= 2mS_-(1+\mathcal{P}_{12})\left\{-\frac{1-x_3}{x_2}G^*+x_1\left[A\left(\frac{\kappa_1^*}{x_1}-\frac{\kappa_2^*}{x_2}\right)+\frac{\kappa^*}{a_1b_2}\right]\right\},\\
\mathcal{M}^{--}_{+-} &= 0,\\
\mathcal{M}^{+-}_{+-} &= 2mS_-\left(F+\mathcal{P}_{12}G\right),
\end{aligned}\tag{42}$$

---

[7] To clarify the notation we stress that in Eqs. (40) and (42) the operator $\mathcal{P}_{12}$ simply changes the indices $1\leftrightarrow 2$, for instance

$$\mathcal{P}_{12}\mathcal{M}^{+-}_{+-}=2mS_-(\mathcal{P}_{12}F+G),\quad \mathcal{P}_{12}F=x_2\left[(\mathcal{P}_{12}A)\frac{\rho_1}{x_1x_3}-\frac{\kappa}{a_2a}\right],$$

with

$$\mathcal{P}_{12}A=\frac{x_3}{a_2a}-\frac{1-x_2}{a_2b_1}+\frac{1}{b_1b}.$$

$$\mathcal{M}^{-+}_{+-} = \mathcal{P}_{12}\mathcal{M}^{+-}_{+-} \,,$$
$$F = x_1\left(A\frac{\rho_2}{x_2x_3} - \frac{\kappa}{a_1 a}\right), \quad G = \frac{x_2}{x_3}\left(A\frac{\kappa_1}{x_1} + \frac{\kappa}{b_2 b}\right).$$

The remaining amplitudes with $\lambda_{e1} = -1/2$ can be obtained by applying the parity conservation relation

$$\mathcal{M}^{-\lambda_1,\,-\lambda_2}_{-\lambda_{e1},-\lambda_{e3}} = -(-1)^{\lambda_{e1}+\lambda_{e3}}\left(\mathcal{M}^{\lambda_1,\,\lambda_2}_{\lambda_{e1},\lambda_{e3}}\right)^* \,. \tag{43}$$

This rule follows from the transformation properties of the electron and photon wave functions under space inversion [see §16 in Ref. [11] and Eq. (25)]:

$$u_i \equiv u^{(\lambda_{ei})}_{\mathbf{p}_i} \to \mathrm{i}\,(-1)^{s-\lambda_{ei}} u^{(-\lambda_{ei})}_{\mathbf{p}_i} \,, \quad \mathbf{pe}^{(\lambda_i)} \to \mathbf{pe}^{(-\lambda_i)} = -\mathbf{pe}^{(\lambda_i)*} \,, \tag{44}$$

where $s = 1/2$ and $\mathbf{p}$ is an arbitrary 2–vector. These properties give an additional factor $(-1)^{n_\gamma+2s-\lambda_{e1}-\lambda_{e3}}$ where $n_\gamma = 2$ is the number of real photons. Therefore, under replacement (43) we have to take the complex conjugates of the original helicity amplitudes with sign plus for the spin non-flip amplitudes (40) and sign minus for the spin flip ones (42).

All helicity amplitudes in Eqs. (40) and (42) are explicitly proportional to κ or to the function A. Therefore, they vanish $\propto |\mathbf{k}|$ in the limit $|\mathbf{k}| \to 0$. It is clear (although not obvious) that also Δ in Eq. (41) vanishes in that limit since both C and D in (40) tend to zero.

We note the explicit Bose symmetry between the two photons in the amplitudes (40) and (42):

$$\mathcal{M}^{\lambda_1\;\lambda_2}_{\lambda_{e1}\lambda_{e3}}(\kappa_1, x_1; \kappa_2, x_2) = \mathcal{M}^{\lambda_2\;\lambda_1}_{\lambda_{e1}\lambda_{e3}}(\kappa_2, x_2; \kappa_1, x_1) \,. \tag{45}$$

Since the spin–flip amplitudes are proportional to the lepton mass they are negligible compared to the spin non–flip ones for not too small scattering angles

$$\frac{m}{x_{1,2,3}E_1} \ll \theta_{1,2,3} \ll 1 \,. \tag{46}$$

We observe that in our kinematics the transitions with the largest change of helicities between initial and final particles $|\lambda_{e1} - \lambda_{e3} - \lambda_1 - \lambda_2| = 3$ are forbidden, in our case the amplitude $\mathcal{M}^{--}_{+-} = 0$. This is in agreement with the vanishing amplitude for single photon bremsstrahlung with the largest change of helicities $|\lambda_{e1} - \lambda_{e3} - \lambda_1| = 2$, see later Eq. (50).

3 Limiting cases of soft and hard final particles

3.1 The soft photon limit

A useful check of the results obtained is the limiting case in which one of the real photons (belonging to the upper block $J_{\rm up}$) becomes soft. For definiteness let us suppose that this is the second photon which means that $x_2 \ll 1$, $k_2 \to 0$, but $\mathbf{k}_2/x_2$ remains finite.

It is known that in this limit the upper block factorizes and can be presented in the form

$$J_{\rm up} = \sqrt{4\pi\alpha}\, J_{\rm up}(e^-_{\lambda_{e1}}\gamma^* \to e^-_{\lambda_{e3}}\gamma_{\lambda_1})\, A_2\,, \tag{47}$$

where

$$A_2 = \left(\frac{p_1}{p_1k_2} - \frac{p_3}{p_3k_2}\right) e_2^* \tag{48}$$

is the factor of the accompanying classical emission of the second photon. This factor is easily transformed to

$$A_2 = \frac{2}{x_2}\left(\frac{\mathbf{k}_2}{a_2} - \frac{\mathbf{r}_2}{b_2}\right)\mathbf{e}^{(\lambda_2)*}\,. \tag{49}$$

The vertex factor $J_{\rm up}(e^-_{\lambda_{e1}}\gamma^* \to e^-_{\lambda_{e3}}\gamma_{\lambda_1})$ can be found in Ref. [1]:

$$\begin{aligned} &J_{\rm up}(e^-_{\lambda_{e1}}\gamma^* \to e^-_{\lambda_{e3}}\gamma_{\lambda_1}) = 8\pi\alpha\,\sqrt{1-x_1}\,{\rm e}^{{\rm i}\,\varphi_3\lambda_{e3}} \\ &\times\left\{\sqrt{2}\,\mathbf{Q}_1\mathbf{e}^{(\lambda_1)*}\,\frac{1-x_1\delta_{\lambda_1,-2\lambda_{e1}}}{x_1}\delta_{\lambda_{e1},\lambda_{e3}} + mR_1\delta_{\lambda_1,2\lambda_{e1}}\delta_{\lambda_{e1},-\lambda_{e3}}\right\}. \end{aligned} \tag{50}$$

Here we use the notations ($\mathbf{q}_1 = \mathbf{k}_1/x_1$)

$$\begin{aligned} \mathbf{Q}_1 &= \frac{\mathbf{k}_1}{a_1} - \frac{\mathbf{r}_1}{x_3b_1} = \frac{\mathbf{q}_1}{m^2+\mathbf{q}_1^2} - \frac{\mathbf{k}+\mathbf{q}_1}{m^2+(\mathbf{k}+\mathbf{q}_1)^2}\,, \\ R_1 &= \frac{x_1}{a_1} - \frac{x_1}{x_3b_1} = \frac{1}{m^2+\mathbf{q}_1^2} - \frac{1}{m^2+(\mathbf{k}+\mathbf{q}_1)^2}\,. \end{aligned} \tag{51}$$

Therefore, in the soft photon limit the spin non–flip and flip amplitudes are

$$\begin{aligned}
\mathcal{M}^{\lambda_1\ \lambda_2}_{\lambda_{e1}\lambda_{e1}} &= 2S_{\lambda_{e1}}\mathbf{Q}_1\mathbf{e}^{(\lambda_1)*}\frac{1-x_1\delta_{\lambda_1,-2\lambda_{e1}}}{x_1}A_2\,, \\
\mathcal{M}^{\lambda_1\ \ \lambda_2}_{\lambda_{e1},-\lambda_{e1}} &= \sqrt{2}m\,S_{-\lambda_{e1}}\,R_1\,\delta_{\lambda_1,2\lambda_{e1}}A_2\,.
\end{aligned} \tag{52}$$

These expressions coincide with those from the amplitudes (40), (42) in the limit of a soft (second) photon when

$$\begin{aligned}
&a\to a_1\,,\quad b\to b_1\,,\quad \rho_1\to\kappa_1+x_1\kappa\,, \\
&A\to -\frac{x_3}{a_1b_2}+\frac{1}{b_1b_2}\,,\quad \mathcal{P}_{12}A\to\frac{x_3}{a_1a_2}-\frac{1}{b_1a_2}\,.
\end{aligned} \tag{53}$$

This can be easily shown for most of the helicity amplitudes given in Eqs. (40), (42), for the amplitude $\mathcal{M}^{+-}_{++}$ we use the form (41) for C and neglect Δ in this limit.

3.2 The soft electron limit $m \ll E_3 \ll E_1$

For this limit we assume that the final electron has an energy E_3 much smaller than that of the initial lepton E_1, but the lepton remains ultrarelativistic. This leads to the condition $x_3 \ll 1$ whereas the value of the 2–vector $\mathbf{p}_3$ is comparable to those of the other 2–vectors. In that limit the helicity amplitudes vanish with $\sqrt{x_3}$ due to the factor $S_{\lambda_{e3}}$ [see Eq. (39)] and the cross section tends to zero with x_3. Besides of this obvious factor the remaining dominant part of the amplitude (staying finite in that limit) can be found.

The kinematic invariants a, b and b_i are simplified to

$$\begin{aligned}
&a=m^2+(\mathbf{k}_1+\mathbf{k}_2)^2\,,\quad x_3b=m^2+(\mathbf{k}+\mathbf{k}_1+\mathbf{k}_2)^2=m^2+\mathbf{p}_3^2\,, \\
&b_i=x_ib\,.
\end{aligned} \tag{54}$$

Additionally we have

$$A\to\frac{x_3}{a_1}\left(\frac{1}{a}-\frac{1}{x_3b}\right)\,,\quad \mathcal{P}_{12}A\to\frac{x_3}{a_2}\left(\frac{1}{a}-\frac{1}{x_3b}\right)\,,\quad \rho_i\to -x_i\pi_3\,. \tag{55}$$

From Eqs. (40) and (42) we derive

$$\begin{aligned}
\mathcal{M}^{++}_{++} &= 2S_+\left(\frac{\kappa_1^*}{x_1a_1}+\frac{\kappa_2^*}{x_2a_2}\right)\left(\frac{\kappa_1^*+\kappa_2^*}{a}+\frac{\pi_3^*}{x_3b}\right)\,, \\
\mathcal{M}^{--}_{++} &= 0\,, \\
\mathcal{M}^{+-}_{++} &= -2x_1S_+\left\{\frac{\kappa_2}{x_2a_2}\left(\frac{\kappa_1^*+\kappa_2^*}{a}+\frac{\pi_3^*}{x_3b}\right)-m^2\frac{1}{a_1}\left(\frac{1}{a}-\frac{1}{x_3b}\right)\right\}\,,
\end{aligned}$$

$$\begin{aligned}
&\mathcal{M}^{-+}_{++} = \mathcal{P}_{12}\mathcal{M}^{+-}_{++}\,, \qquad\qquad (56)\\
&\mathcal{M}^{++}_{+-} = -2mS_-\left(\frac{1}{a}-\frac{1}{x_3 b}\right)\left(\frac{\kappa_1^*}{x_1 a_1}+\frac{\kappa_2^*}{x_2 a_2}\right)\,,\\
&\mathcal{M}^{+-}_{+-} = 2mx_1S_-\left\{\frac{1}{a_1}\left(\frac{\kappa_1+\kappa_2}{a}+\frac{\pi_3}{x_3 b}\right)+\left(\frac{1}{a}-\frac{1}{x_3 b}\right)\frac{\kappa_2}{x_2 a_2}\right\}\,,\\
&\mathcal{M}^{-+}_{+-} = \mathcal{P}_{12}\mathcal{M}^{+-}_{+-}\,,\\
&\mathcal{M}^{--}_{+-} = 0\,.
\end{aligned}$$

Introducing the notations

$$\mathbf{Q}_e = \frac{\mathbf{k}_1+\mathbf{k}_2}{a}+\frac{\mathbf{p}_3}{x_3 b}\,, \quad R_e = \frac{1}{a}-\frac{1}{x_3 b}\,, \qquad (57)$$

the matrix elements in the soft electron limit can be written as

$$\begin{aligned}
\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}} &= \sqrt{x_3}s\,\mathrm{e}^{\mathrm{i}\,\varphi_3\lambda_{e3}}\,(1+\mathcal{P}_{12})\left(2\mathbf{Q}_e\mathbf{e}^{(\lambda_2)*}\,\delta_{\lambda_2,2\lambda_{e3}}+\sqrt{2}mR_e\delta_{\lambda_2,-2\lambda_{e3}}\right)\\
&\times\frac{1}{a_1}\left(\frac{1-x_1\delta_{\lambda_1,-\lambda_2}}{x_1}2\mathbf{k}_1\mathbf{e}^{(\lambda_1)*}\,\delta_{\lambda_2,2\lambda_{e1}}+\sqrt{2}mx_1\delta_{\lambda_1,-\lambda_2}\delta_{\lambda_2,-2\lambda_{e1}}\right)\,. \qquad (58)
\end{aligned}$$

Considering additionally a soft (second) photon we arrive at the case (60) discussed in Section 3.3.

3.3 The limits of a hard electron or a hard photon

In the limit of a hard electron $x_3 \to 1$ both photons become soft. Using Eqs. (52) for that case, only the non–flip amplitudes remain:

$$\begin{aligned}
\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}} &= s\,A_1A_2\,\mathrm{e}^{\mathrm{i}\,\varphi_3\lambda_{e3}}\delta_{\lambda_{e1},\lambda_{e3}}\,, \qquad (59)\\
A_i = \frac{2}{x_i}\left(\frac{\mathbf{k}_i}{a_i}-\frac{\mathbf{r}_i}{b_i}\right)\mathbf{e}^{(\lambda_i)*} &= 2\left(\frac{\mathbf{k}_i\mathbf{e}^{(\lambda_i)*}}{x_i^2m^2+\mathbf{k}_i^2}-\frac{(\mathbf{k}_i+x_i\mathbf{k})\mathbf{e}^{(\lambda_i)*}}{x_i^2m^2+(\mathbf{k}_i+x_i\mathbf{k})^2}\right)\,.
\end{aligned}$$

Now we consider the case when the first photon is hard and the soft electron remains ultrarelativistic: $x_1 \to 1$ leading to $x_{2,3} \ll 1$, $|\mathbf{r}_2|/b_2 \ll |\mathbf{k}_2|/a_2$ and $A_2 \to 2\mathbf{k}_2\mathbf{e}^{(\lambda_2)*}/(x_2a_2)$. The amplitudes can be derived from Eqs. (52) where $\mathbf{Q}_1$ and R_1 from Eqs. (51) have to be taken at $x_1 \to 1$:

$$\mathcal{M}^{\lambda_1\,\lambda_2}_{\lambda_{e1}\lambda_{e3}} = \sqrt{x_3}s\,\mathrm{e}^{\mathrm{i}\,\varphi_3\lambda_{e3}}\left(2\mathbf{Q}_1\mathbf{e}^{(\lambda_1)*}\delta_{\lambda_{e3},\lambda_{e1}}+\sqrt{2}m\,R_1\delta_{-\lambda_{e3},\lambda_{e1}}\right) \qquad (60)$$

$$\times\delta_{\lambda_1,2\lambda_{e1}}\,\frac{2}{x_2 a_2}\mathbf{k}_2\mathbf{e}^{(\lambda_2)*}\,.$$

From that expression one can see that only those amplitudes remain in which the initial electron transfers its helicity to the hard photon: $\lambda_1 = 2\lambda_{e1}$, and they vanish as $\sqrt{x_3}$.

4 The cross channel $\gamma e^\pm \to \gamma e^+ e^- + e^\pm$

4.1 Crossing relations and helicity amplitudes

We consider the reaction

$$\gamma(\tilde{k}_1) + e^\pm(p_2) \to \gamma(\tilde{k}_2) + e^+(p_+) + e^-(p_-) + e^\pm(p_4)\,, \tag{61}$$

which is one of the cross channels to the double forward bremsstrahlung considered before. We denote in the cross channel the 4–momenta of the initial photon by $\tilde{k}_1$ with energy $\tilde{\omega}_1$ and helicity λ_1, the 4–momenta (Euclidean 2–vectors) of the final lepton pair by $p_\pm$ ($\mathbf{p}_\pm$) and the final photon by $\tilde{k}_2$ ($\tilde{\mathbf{k}}_2$) with helicities $\lambda_\pm$ and λ_2, respectively.

The amplitude for reaction (61) can be obtained in the jet kinematics by considering the cross channel of the upper vertex only:

$$M(\gamma e^\pm \to \gamma e^+ e^- + e^\pm) = \frac{\tilde{s}}{k^2} J_{\rm up}(\gamma\gamma^* \to \gamma e^+ e^-) J_{\rm down}(e^\pm\gamma^* \to e^\pm)\,, \tag{62}$$

with

$$J_{\rm up}(\gamma\gamma^* \to \gamma e^+ e^-) = J_{\rm up}^{\rm cross}(e^-\gamma^* \to e^-\gamma\gamma)\,, \quad \tilde{s} = 2\tilde{k}_1 p_2\,. \tag{63}$$

The following 4–vector replacements have to be performed:

$$k_1 \to -\tilde{k}_1\,, \quad k_2 \to \tilde{k}_2\,, \quad p_1 \to -p_+\,, \quad p_3 \to p_-\,, \tag{64}$$

with the remaining 4–vectors k, p_2 and p_4 held intact.

Obviously, the corresponding energy fractions are

$$y_\pm = \frac{E_\pm}{\tilde{\omega}_1}\,, \quad y_2 = \frac{\tilde{\omega}_2}{\tilde{\omega}_1}\,, \quad y_+ + y_- + y_2 = 1\,. \tag{65}$$

The Sudakov decomposition is easily performed (using $\tilde{k}_1$ and $\tilde{p}' = p_2 - (m^2/\tilde{s})\tilde{k}_1$ as light–like vectors instead of p and p', respectively)

$$p_\pm = \frac{m^2 + \mathbf{p}_\pm^2}{\tilde{s} y_\pm}\,\tilde{p}' + y_\pm \tilde{k}_1 + p_{\pm\perp}\,, \quad \tilde{k}_2 = \frac{\tilde{\mathbf{k}}_2^2}{\tilde{s} y_2}\,\tilde{p}' + y_2 \tilde{k}_1 + \tilde{k}_{2\perp}\,. \tag{66}$$

Comparing the kinematic invariants a_i, b_i with their corresponding crossed invariants and using the replacements (64) and the Sudakov decomposition (66) we find the substitution rules for the energy fractions and Euclidean 2–vectors (compare Ref. [12])

$$\begin{aligned} & x_1 \to \frac{1}{y_+}\,, \quad x_2 \to -\frac{y_2}{y_+}\,, \quad x_3 \to -\frac{y_-}{y_+}\,, \\ & \frac{\mathbf{k}_1}{x_1} \to \mathbf{p}_+\,, \quad \frac{\mathbf{r}_1}{x_1} \to -\mathbf{p}_-\,, \quad \frac{\mathbf{k}_2}{x_2} \to \mathbf{p}_+ - \frac{y_+}{y_2}\tilde{\mathbf{k}}_2\,, \quad \frac{\mathbf{r}_2}{x_2} \to -\mathbf{p}_- + \frac{y_-}{y_2}\tilde{\mathbf{k}}_2\,, \end{aligned} \tag{67}$$

with

$$\mathbf{p}_+ + \mathbf{p}_- + \tilde{\mathbf{k}}_2 + \mathbf{k} = 0\,.$$

To take into account the substitution rules for the polarizations of the particles, we have to re-introduce the azimuthal angle of the initial electron φ_1. This changes only the phase of the factor $S_{\lambda_{e3}}$ given in Eq. (39). Therefore, in this section we use

$$\begin{aligned} S_{\lambda_{e1}\lambda_{e3}}(\varphi_1,\varphi_3) &= s\sqrt{x_3}\,\mathrm{e}^{\mathrm{i}\,(\varphi_3\lambda_{e3}-\varphi_1\lambda_{e1})} = 4E_1E_2\sqrt{x_3}\,\mathrm{e}^{\mathrm{i}\,(\varphi_3\lambda_{e3}-\varphi_1\lambda_{e1})}\,, \\ S_{\lambda_{e3}} &= S_{\lambda_{e1}\lambda_{e3}}(0,\varphi_3) \end{aligned} \tag{68}$$

instead of $S_{\lambda_{e3}}$. Consequently, the helicity amplitudes given in Eqs. (40) and (42) have to be used now with the factor $S_{\lambda_{e1}\lambda_{e3}}(\varphi_1,\varphi_3)$ and they depend on both lepton azimuthal angles explicitly.

Under crossing we have the substitution rules for the helicities and the lepton azimuthal angles

$$\lambda_1 \to -\lambda_1\,, \quad \lambda_{e1} \to -\lambda_+\,, \quad \lambda_{e3} \to \lambda_-\,, \quad \lambda_2 \to \lambda_2\,, \quad \varphi_{1,3} \to \varphi_{+,-}\,. \tag{69}$$

which leads to

$$S_{\lambda_{e1}\lambda_{e3}}(\varphi_1,\varphi_3) \to \widetilde{S}_{\lambda_+\lambda_-} = -\mathrm{i}\,\tilde{s}\sqrt{y_+y_-}\,\mathrm{e}^{\mathrm{i}\,(\varphi_-\lambda_-+\varphi_+\lambda_+)}\,. \tag{70}$$

The kinematic invariants are transformed to

$$\begin{aligned}
&\left\{\begin{matrix}a_1\\ b_1\end{matrix}\right\} \to a_\pm = \pm 2\tilde{k}_1 p_\pm = \pm\frac{1}{y_\pm}\left(m^2+\mathbf{p}_\pm^2\right)\,,\\
&\left\{\begin{matrix}a_2\\ b_2\end{matrix}\right\} \to b_\pm = \mp 2\tilde{k}_2 p_\pm = \mp\frac{1}{y_\pm y_2}\left[y_2^2 m^2+\left(y_2\mathbf{p}_\pm - y_\pm\tilde{\mathbf{k}}_2\right)^2\right]\,,\\
&\left\{\begin{matrix}a\\ b\end{matrix}\right\} \to \tilde{a}_\pm = \mp k^2 \mp 2kp_\mp = a_\pm + b_\pm \pm \frac{1}{y_2}\tilde{\mathbf{k}}_2^2\,,\\
&A \to y_- B = y_-\left\{-\frac{1}{y_+a_+\ \tilde{a}_+}+\frac{1-y_+}{y_+a_+\ y_-b_-}+\frac{1}{y_-b_-\ \tilde{a}_-}\right\}\,.
\end{aligned} \tag{71}$$

We introduce the permutation operator $\mathcal{P}$ with the properties

$$\mathcal{P}f(p_+,\lambda_+;p_-,\lambda_-) = f(p_-,\lambda_-;p_+,\lambda_+)\,, \tag{72}$$

or in more detail

$$\mathcal{P}f(a_\pm,b_\pm,\tilde{a}_\pm,y_\pm,\pi_\pm,\lambda_\pm) = f(-a_\mp,-b_\mp,-\tilde{a}_\mp,y_\mp,\pi_\mp,\lambda_\mp)\,. \tag{73}$$

As usual, the circular components of $\mathbf{p}_\pm$ and $\tilde{\mathbf{k}}_2$ are used

$$\pi_\pm = p_{x\pm} + \mathrm{i}\,p_{y\pm}\,, \quad \tilde{\kappa}_2 = \tilde{k}_{2x} + \mathrm{i}\,\tilde{k}_{2y}\,. \tag{74}$$

The operator $\mathcal{P}$ is similar to $\mathcal{P}_{12}$ in the original channel, in fact it will interchange $e^+ \leftrightarrow e^-$ in the crossed amplitudes.

As a result, we obtain the amplitude for reaction (61) in the considered jet kinematics

$$M(\gamma e^\pm \to \gamma e^+e^- + e^\pm) = \frac{32\pi^2\alpha^2}{k^2}\widetilde{\mathcal{M}}^{\lambda_1\,\lambda_2}_{\lambda_+\lambda_-}\,\mathrm{e}^{-\mathrm{i}\,\lambda_{e4}\,\varphi_4}\,\delta_{\lambda_{e2},\lambda_{e4}}\,, \tag{75}$$

where λ_{e2} (λ_{e4}) are the helicities of the electrons or positrons moving along the $-z$ axis before and after the collision with the initial photon. Using the function C in Eq. (41) for crossing and the notations

$$Y_1 = B\pi_+ + \frac{\kappa}{y_-b_-\,\tilde{a}_-}\,, \quad Y_2 = \pi_- - \frac{y_-}{y_2}\tilde{\kappa}_2\,, \tag{76}$$

the helicity amplitudes in the cross channel are presented as follows:

$$\widetilde{\mathcal{M}}^{-+}_{-+} = 2\widetilde{S}_{-+}\,y_+(1-\mathcal{P})\left\{Y_1^*Y_2^* - \frac{\pi_+^*\kappa^*}{y_+a_+\,\tilde{a}_+}\right\}\,,$$

$$
\begin{aligned}
\widetilde{\mathcal{M}}^{+-}_{-+} &= \widetilde{S}_{-+}\mathcal{P}\left(\frac{\widetilde{\mathcal{M}}^{-+}_{-+}}{\widetilde{S}_{-+}}\right)^{*} , \\
\widetilde{\mathcal{M}}^{--}_{-+} &= -2\widetilde{S}_{-+}\left(\widetilde{C}+\mathcal{P}\widetilde{D}^{*}\right) , \\
\widetilde{\mathcal{M}}^{++}_{-+} &= \widetilde{S}_{-+}\mathcal{P}\left(\frac{\widetilde{\mathcal{M}}^{--}_{-+}}{\widetilde{S}_{-+}}\right)^{*} , \\
\widetilde{D} &= -(1-y_{+})\left\{Y_1 Y_2^{*} - \frac{\pi_{+}\kappa^{*}}{y_{+}a_{+}\,\tilde{a}_{+}}\right\} , \\
\widetilde{C} &= -\frac{1-y_{-}}{1-y_{+}}\widetilde{D}^{*} - \frac{1}{\tilde{a}_{+}} + \frac{1}{\tilde{a}_{-}} + \frac{y_2\left(m^2 - Y_2\pi_{+}^{*}\right)}{y_{+}a_{+}\,y_{-}b_{-}} - \frac{\pi_{+}^{*}\tilde{\kappa}_2}{y_{+}a_{+}\,\tilde{a}_{+}} + \frac{\tilde{\kappa}_2^{*}Y_2}{y_{-}b_{-}\,\tilde{a}_{-}} , \\
\widetilde{\mathcal{M}}^{--}_{--} &= 2m\widetilde{S}_{--}\,(1+\mathcal{P})\left\{BY_2 - \frac{\kappa}{y_{+}a_{+}\,\tilde{a}_{+}}\right\} , \\
\widetilde{\mathcal{M}}^{++}_{--} &= 2m\widetilde{S}_{--}\,y_2\,(1+\mathcal{P})Y_1 , \\
\widetilde{\mathcal{M}}^{-+}_{--} &= 2m\widetilde{S}_{--}\,(1+\mathcal{P})\left\{(1-y_2)Y_1^{*} + B\frac{y_{-}}{y_2}\tilde{\kappa}_2^{*} + \frac{\kappa^{*}}{y_{+}a_{+}\,b_{-}}\right\} , \\
\widetilde{\mathcal{M}}^{+-}_{--} &= 0 .
\end{aligned}
\tag{77}
$$

The remaining amplitudes are derived from a relation equivalent to Eq. (43)

$$
\widetilde{\mathcal{M}}^{-\lambda_1,\,-\lambda_2}_{-\lambda_{+},-\lambda_{-}} = -(-1)^{\lambda_{+}+\lambda_{-}}\left(\widetilde{\mathcal{M}}^{\lambda_1,\,\lambda_2}_{\lambda_{+},\lambda_{-}}\right)^{*} . \tag{78}
$$

The helicity amplitudes (77), (78) are C–odd. Therefore, they have to be symmetric under lepton–antilepton exchange $e^{-} \leftrightarrow e^{+}$. This symmetry, expressed by the relation

$$
\begin{aligned}
\widetilde{\mathcal{M}}^{\lambda_1\,\lambda_2}_{\lambda_{+}\lambda_{-}}\,(a_{\pm}, b_{\pm}, \tilde{a}_{\pm}, y_{\pm}, \pi_{\pm}, \varphi_{\pm}) = \\
\widetilde{\mathcal{M}}^{\lambda_1\,\lambda_2}_{\lambda_{-}\lambda_{+}}\,(-a_{\mp}, -b_{\mp}, -\tilde{a}_{\mp}, y_{\mp}, \pi_{\mp}, \varphi_{\mp}) ,
\end{aligned}
\tag{79}
$$

can be easily seen for all amplitudes taking into account Eq. (78) and $\mathcal{P}^2 = 1$.

Note an alternative form for $\widetilde{C}$ useful for deriving the hard photon limit discussed below in Section 4.2:

$$
\widetilde{C} = \frac{y_{-}}{1-y_{+}}\widetilde{D}^{*} - B\left(\frac{y_{-}\tilde{\kappa}_2\pi_{+}^{*}}{y_2} - m^2\right) - \frac{\tilde{\kappa}_2\kappa^{*}}{y_2 b_{-}\tilde{a}_{-}} - \frac{\kappa\pi_{+}^{*}}{y_{+}a_{+}b_{-}} . \tag{80}
$$

Since the function B analogous to A tends to zero for $|\mathbf{k}| \to 0$, the helicity amplitudes (77) are again proportional to $|\mathbf{k}|$ in that limit as expected from crossing. For $\widetilde{C}$ this can be seen using its form (80).

4.2 Soft and hard particle limits

In the soft photon limit ($\tilde{k}_2 \to 0$) we use

$$\mathbf{Q} = \frac{\mathbf{p}_+}{m^2+\mathbf{p}_+^2} + \frac{\mathbf{p}_-}{m^2+\mathbf{p}_-^2}\,, \qquad R = \frac{1}{m^2+\mathbf{p}_+^2} - \frac{1}{m^2+\mathbf{p}_-^2} \tag{81}$$

and have to consider (at $y_2 \to 0$)

$$\tilde{a}_\pm \to a_\pm\,, \quad B \to \frac{R}{b_-}\,, \quad \mathcal{P}B \to \frac{R}{b_+}\,. \tag{82}$$

This limit again leads to a factorized form of the upper vertex:

$$J_{\mathrm{up}}(\gamma_{\lambda_1}\gamma^* \to e^+_{\lambda_+}e^-_{\lambda_-}\gamma_{\lambda_2}) = \sqrt{4\pi\alpha}\,J_{\mathrm{up}}(\gamma_{\lambda_1}\gamma^* \to e^+_{\lambda_+}e^-_{\lambda_-})\,\widetilde{A}_2\,, \tag{83}$$

where $J_{\mathrm{up}}(\gamma_{\lambda_1}\gamma^* \to e^+_{\lambda_+}e^-_{\lambda_-})$ can be found in Ref. [1] and

$$\widetilde{A}_2 = \left(\frac{p_+}{p_+\tilde{k}_2} - \frac{p_-}{p_-\tilde{k}_2}\right)e_2^* = \frac{2}{y_2}\left(\frac{y_2\mathbf{p}_+ - y_+\tilde{\mathbf{k}}_2}{b_+} + \frac{y_2\mathbf{p}_- - y_-\tilde{\mathbf{k}}_2}{b_-}\right)\mathbf{e}^{(\lambda_2)*} \tag{84}$$

is the classical emission factor of the final photon. With notations (81) and (84) we find the non-vanishing helicity amplitudes in that limit

$$\begin{aligned}\widetilde{\mathcal{M}}^{\lambda_1\,\lambda_2}_{\lambda_+\lambda_-} =\,& \mathrm{i}\,\tilde{s}\,\sqrt{y_+y_-}\mathrm{e}^{\mathrm{i}\,(\varphi_+\lambda_++\varphi_-\lambda_-)}\widetilde{A}_2 \\ &\times\left\{2\mathbf{Q}\mathbf{e}^{(\lambda_1)}\,\left(y_+ - \delta_{\lambda_1,-2\lambda_+}\right)\delta_{\lambda_+,-\lambda_-} - \sqrt{2}mR\delta_{\lambda_1,2\lambda_+}\delta_{\lambda_+,\lambda_-}\right\}.\end{aligned} \tag{85}$$

From Eq. (85) we derive the amplitudes in the limit of a hard electron ($y_- \to 1$, $y_+, y_2 \ll 1$)

$$\begin{aligned}\widetilde{\mathcal{M}}^{\lambda_1\,\lambda_2}_{\lambda_+\lambda_-} =\,& -\mathrm{i}\,\tilde{s}\,\sqrt{y_+}\mathrm{e}^{\mathrm{i}\,(\varphi_+\lambda_++\varphi_-\lambda_-)}\frac{2}{b_-}\mathbf{p}_-\mathbf{e}^{(\lambda_2)*} \\ &\times\left\{2\mathbf{Q}\mathbf{e}^{(\lambda_1)}\,\delta_{\lambda_+,-\lambda_-} + \sqrt{2}mR\delta_{\lambda_+,\lambda_-}\right\}\delta_{\lambda_1,2\lambda_-}\,.\end{aligned} \tag{86}$$

In this case the helicity of the initial photon is transferred to that of the hard electron only, i.e., $\lambda_1 = 2\lambda_-$.

The soft electron limit corresponds to the condition $y_- \ll 1$ and finite transverse momentum $\mathbf{p}_-$. It can be obtained as cross channel from Eq. (58) performing the replacements (69) and (70) for $S_{\lambda_{e1}\lambda_{e3}}$ as well as

$$\begin{aligned}
&x_1 \to \frac{1}{y_+}\,, \quad x_2 \to -\frac{y_2}{y_+} = -\frac{1-y_+}{y_+}\,, \\
&\mathbf{k}_1 \to \frac{1}{y_+}\mathbf{p}_+\,, \quad \mathbf{k}_2 \to -\frac{1-y_+}{y_+}\mathbf{p}_+ + \tilde{\mathbf{k}}_2\,, \quad \mathbf{p}_3 \to \mathbf{p}_-\,, \\
&a_1 \to a_+\,, \quad a_2 \to b_+\,, \quad a \to m^2 + (\mathbf{p}_+ + \tilde{\mathbf{k}}_2)^2\,, \quad x_3 b \to m^2 + \mathbf{p}_-^2\,.
\end{aligned} \tag{87}$$

In that limit the helicity amplitudes vanish as $\sqrt{y_-}$.

Finally we consider the hard final photon limit $y_2 \to 1$ and $y_\pm \ll 1$. From the general helicity amplitudes (77) and using the form (80) of $\widetilde{C}$ we find in leading order in y_+ and y_-

$$\begin{aligned}
\widetilde{\mathcal{M}}^{--}_{-+} = \widetilde{\mathcal{M}}^{++}_{-+} &= -2\widetilde{S}_{-+}\left\{B\left(m^2 - \pi_-^*\pi_+\right) + \frac{\kappa^*\pi_+}{y_+a_+\,\tilde{a}_+} + \frac{\pi_-^*\kappa}{y_-a_-\,\tilde{a}_-}\right\}, \\
\widetilde{\mathcal{M}}^{++}_{--} = \widetilde{\mathcal{M}}^{--}_{--} &= 2m\widetilde{S}_{--}(1+\mathcal{P})\left\{B\pi_- - \frac{\kappa}{y_+a_+\,\tilde{a}_+}\right\}, \\
\widetilde{\mathcal{M}}^{+\,-}_{\lambda_+\lambda_-} = \widetilde{\mathcal{M}}^{-\,+}_{\lambda_+\lambda_-} &= 0\,,
\end{aligned} \tag{88}$$

where in this limit $b_\pm \to -a_\pm$ and

$$\begin{aligned}
B = \mathcal{P}B &= -\frac{1}{y_+a_+\,\tilde{a}_+} - \frac{1}{y_+a_+\,y_-a_-} - \frac{1}{y_-a_-\,\tilde{a}_-}\,, \\
\tilde{a}_+ = -\mathcal{P}\tilde{a}_- &= y_-a_+ + m^2 + (\mathbf{p}_+ + \tilde{\mathbf{k}}_2)^2\,,
\end{aligned} \tag{89}$$

the $a_\pm$ are defined in Eqs. (71). Note that the helicity of the initial photon is transferred to that of the hard final photon, the amplitudes vanish as $\sqrt{y_+y_-}$.

5 Conclusions

The present work completes a series of publications [2,3,1] where the helicity analysis of Born processes with non-decreasing cross sections up to fourth order in α has been performed in a jet kinematics. These specific kinematic conditions provide the main contribution to the total cross section at high energies. On the other hand, under these kinematic conditions Monte Carlo simulations as well as analytical calculations by known methods (compare, for example, Ref. [9]) are difficult to perform. Therefore, the results obtained can be useful for monitoring and calibrating of polarized beams at high energy colliders. They might also be used for almost exact estimates of the essential background in a number of reactions with polarized particles.

The main results of our paper are summarized in Eqs. (37), (40), (42) and (75), (77) which give the analytical expressions for all 64 helicity amplitudes to the high accuracy (3). Let us summarize some specific features of those helicity amplitudes using as example the double bremsstrahlung process:

- The obtained formulae for the helicity amplitudes are compact and convenient for numeric calculations since in their form large compensating terms are already cancelled. Indeed, in Eqs. (40) and (42) all items are either proportional to $\kappa = k_x + \mathrm{i}\, k_y$ or to the function A (34) which vanishes at small transverse momentum of the t–channel exchange photon $|\mathbf{k}|$. Therefore, the amplitude

$$|M(e^- e^+ \to e^- \gamma\gamma + e^+)| \propto |\mathbf{k}| \;\text{ at }\; |\mathbf{k}| \to 0\,. \tag{90}$$

- The helicity amplitudes $\mathcal{M}_{+\,-}^{\lambda_1\lambda_2}$ and $\mathcal{M}_{-\,+}^{\lambda_1\lambda_2}$, in which the electron changes its helicity, are proportional to the electron mass m (see Eq. (42)). They become small compared with the non-spin flip amplitudes $\mathcal{M}_{+\,+}^{\lambda_1\lambda_2}$ and $\mathcal{M}_{-\,-}^{\lambda_1\lambda_2}$ for large scattering angles (46) or small energies of both photons $\omega_{1,2} \ll E_1$.
- The amplitudes $\mathcal{M}_{+\,-}^{-\,-}$ and $\mathcal{M}_{-\,+}^{+\,+}$ with the maximal change of helicities $|\lambda_{e1} - \lambda_{e3} - \lambda_1 - \lambda_2| = 3$ are equal to zero under our kinematic conditions.
- If the energy of any final particle tends to that of the initial electron, the incoming electron transfers its helicity to that particle: $\lambda_i = 2\lambda_{e1}$ at $\omega_i \to E_1$ or $\lambda_{e3} = \lambda_{e1}$ at $E_3 \to E_1$.

Acknowledgements

This work is supported in part by Volkswagen Stiftung (Az. No. I/72 302) and by Russian Foundation for Basic Research (codes 96-15-96030, 99-02-17211 and 99-02-17730).

A Calculation of necessary bilinear spinor structures

We have to calculate all bilinear spinor structures appearing in Eq. (36) in order to find the helicity amplitudes in the $s \to \infty$ limit. For example, let us consider the numerator of the second term of Eq. (36).

$$N = \bar{u}_3 \hat{p}' \hat{k}_\perp \hat{e}_2^* (\hat{p}_1 - \hat{k}_1 + m) \hat{e}_1^* u_1\,. \tag{A.1}$$

We transpose $\hat{e}_1^*$ to the left and obtain

$$N = \bar{u}_3 \hat{p}' \hat{k}_\perp \hat{e}_2^* \left[\hat{e}_1^* (-\hat{p}_1 + \hat{k}_1 + m) + 2 p_1 e_1^* \right] u_1 \, . \tag{A.2}$$

Using the Sudakov decomposition of the 4–vectors e_i^* and k_1 [see Eqs. (13), (24), (26)] and taking into account $\hat{p}'\hat{p}' = 0$, $\hat{p} u_1 \approx \hat{p}_1 u_1 = m u_1$ we find

$$N = \bar{u}_3 \hat{p}' \hat{k}_\perp \hat{e}_{2\perp}^* \left[\hat{e}_{1\perp}^* \hat{k}_{1\perp} + m x_1 \hat{e}_{1\perp}^* + \frac{2}{x_1} \mathbf{k}_1 \mathbf{e}^{(\lambda_1)*} \right] u_1 \, . \tag{A.3}$$

Similar transformations can be performed for the other terms of Eq. (36).

As a result we conclude that it is sufficient to calculate the following combinations $\bar{u}_3 \hat{p}' \hat{a}_\perp \hat{b}_\perp \ldots \hat{c}_\perp u_1$ and $\bar{u}_3 \hat{a}_\perp \hat{b}_\perp \ldots \hat{c}_\perp u_1$. Here the bispinors u are defined as follows (for an initial electron with helicity +1/2)

$$u_1 = \begin{pmatrix} \sqrt{E_1 + m}\, w_1 \\ \sqrt{E_1 - m}\, w_1 \end{pmatrix} , \quad u_3 = \begin{pmatrix} \sqrt{E_3 + m}\, w_3 \\ 2\lambda_{e3} \sqrt{E_3 - m}\, w_3 \end{pmatrix} , \tag{A.4}$$

with

$$w_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} , \quad w_3^{(+1/2)} = \begin{pmatrix} \mathrm{e}^{-\mathrm{i}\,\varphi_3/2} \\ \frac{\theta_3}{2}\, \mathrm{e}^{\mathrm{i}\varphi_3/2} \end{pmatrix} , \quad w_3^{(-1/2)} = \begin{pmatrix} -\frac{\theta_3}{2}\, \mathrm{e}^{-\mathrm{i}\varphi_3/2} \\ \mathrm{e}^{\mathrm{i}\varphi_3/2} \end{pmatrix} , \tag{A.5}$$

and

$$\hat{p}' = E_2 \begin{pmatrix} 1 & \sigma_z \\ -\sigma_z & -1 \end{pmatrix} , \quad \hat{a}_\perp = -a_x \gamma_x - a_y \gamma_y \, . \tag{A.6}$$

The σ_i and γ_i are the Pauli and Dirac matrices in the standard representation. Introducing the circular components of the Euclidean 2–vector $\mathbf{a}$

$$a_\pm = a_x \pm \mathrm{i}\, a_y \, , \qquad a_\mp^* = a_\pm \, , \tag{A.7}$$

we obtain

$$\hat{a}_\perp = -\frac{1}{2} (a_+ \gamma_- + a_- \gamma_+) \, , \quad \gamma_\pm = \begin{pmatrix} 0 & \sigma_\pm \\ -\sigma_\pm & 0 \end{pmatrix} \, . \tag{A.8}$$

Since $\sigma_+ w_1 = 0$ and $\sigma_- w_1 = \binom{0}{2}$, we have $\gamma_+ u_1 = 0$ and $\gamma_+\gamma_- u_1 = -4u_1$. From this it follows that

$$\hat{a}_\perp u_1 = -\frac{1}{2} a_+ \gamma_- u_1 \,, \quad \hat{a}_\perp \hat{b}_\perp u_1 = -a_- b_+ u_1 \,, \qquad \text{(A.9)}$$
$$\hat{a}_\perp \hat{b}_\perp \hat{c}_\perp u_1 = \frac{1}{2} a_+ b_- c_+ \gamma_- u_1 \,, \quad \hat{a}_\perp \hat{b}_\perp \hat{c}_\perp \hat{d}_\perp u_1 = a_- b_+ c_- d_+ u_1 \,.$$

Introducing the shorthand notations

$$\delta_{\lambda_{e3},\pm 1/2} = \delta_\pm \,, \quad \delta_{\lambda_i,\pm 1} = \delta_{i\pm}$$

and using the quantity $S_{\lambda_{e3}}$ defined in Eq. (39) we arrive at the following four basic combinations

$$\begin{aligned}
\bar{u}_3 \hat{p}' u_1 &= S_+ \delta_+ \,, \quad \bar{u}_3 \hat{p}' \hat{a}_\perp u_1 = -a_+ S_- \delta_- \,, \\
s\, \bar{u}_3 u_1 &= \frac{1}{x_3} \left[-\pi_3 S_- \delta_- + (1+x_3) m S_+ \delta_+ \right] \,, \\
s\, \bar{u}_3 \hat{a}_\perp u_1 &= -\frac{a_+}{x_3} \left[\pi_3^* S_+ \delta_+ + (1-x_3) m S_- \delta_- \right] \,,
\end{aligned} \qquad \text{(A.10)}$$

where

$$\pi_3 = p_{3x} + \mathrm{i}\, p_{3y} = -\kappa - \kappa_1 - \kappa_2 \,. \qquad \text{(A.11)}$$

Choosing $a_\perp = e_\perp^{(\lambda)*}$ we find $a_\pm = \mp\sqrt{2}\delta_{\lambda,\pm 1}$.

Then the relevant bilinear combinations [using the Sudakov decomposition of all 4–vectors of Eq. (36)] are found to be

$$\begin{aligned}
\bar{u}_3 \hat{p}' u_1 &= S_+ \delta_+ \,, \\
\bar{u}_3 \hat{p}' \hat{e}_i^* u_1 &= \sqrt{2} \delta_{i+} S_- \delta_- \,, \\
\bar{u}_3 \hat{p}' \hat{e}_2^* \hat{e}_1^* u_1 &= 2 \delta_{2-} \delta_{1+} S_+ \delta_+ \,, \\
\bar{u}_3 \hat{p}' \hat{e}_{2\perp}^* \hat{k}_{i\perp} u_1 &= -\sqrt{2} \kappa_i \delta_{2-} S_+ \delta_+ \,, \\
\bar{u}_3 \hat{p}' \hat{k}_\perp \hat{e}_{i\perp}^* u_1 &= \sqrt{2} \kappa^* \delta_{i+} S_+ \delta_+ \,, \\
\bar{u}_3 \hat{p}' \hat{k}_\perp \hat{e}_{2\perp}^* \hat{e}_{1\perp}^* u_1 &= -2\kappa \delta_{2-} \delta_{1+} S_- \delta_- \,, \\
\bar{u}_3 \hat{p}' \hat{e}_{2\perp}^* \hat{e}_{1\perp}^* \hat{k}_{i\perp} u_1 &= -2\kappa_i \delta_{2+} \delta_{1-} S_- \delta_- \,, \\
\bar{u}_3 \hat{p}' \hat{e}_{2\perp}^* \hat{k}_\perp \hat{e}_{1\perp}^* u_1 &= 2\kappa^* \delta_{2+} \delta_{1+} S_- \delta_- \,, \\
\bar{u}_3 \hat{p}' \hat{e}_{2\perp}^* \hat{e}_{1\perp}^* \hat{k}_\perp u_1 &= -2\kappa \delta_{2+} \delta_{1-} S_- \delta_- \,, \\
\bar{u}_3 \hat{p}' \hat{e}_{2\perp}^* \hat{k}_\perp \hat{e}_{1\perp}^* \hat{k}_{i\perp} u_1 &= 2\kappa\kappa_i \delta_{2-} \delta_{1-} S_+ \delta_+ \,, \\
\bar{u}_3 \hat{p}' \hat{k}_{i\perp} \hat{e}_{2\perp}^* \hat{e}_{1\perp}^* \hat{k}_\perp u_1 &= -2\kappa_i^* \kappa \delta_{2+} \delta_{1-} S_+ \delta_+ \,,
\end{aligned} \qquad \text{(A.12)}$$

$$\begin{aligned}
s\,\bar u_3 \hat e^*_{i\perp} u_1 &= \sqrt{2}\delta_{i+}\frac{1}{x_3}\left[\pi_3^* S_+\delta_+ + (1-x_3) m S_-\delta_-\right] \ , \\
s\,\bar u_3 \hat e^*_{2\perp}\hat e^*_{1\perp} u_1 &= 2\delta_{2-}\delta_{1+}\frac{1}{x_3}\left[-\pi_3 S_-\delta_- + (1+x_3) m S_+\delta_+\right] \ , \\
s\,\bar u_3 \hat e^*_{2\perp}\hat e^*_{1\perp}\hat k_{i\perp} u_1 &= -2\delta_{2+}\delta_{1-}\frac{\kappa_i}{x_3}\left[\pi_3^* S_+\delta_+ + (1-x_3) m S_-\delta_-\right] \ , \\
s\,\bar u_3 \hat e^*_{2\perp}\hat k_{\perp}\hat e^*_{1\perp} u_1 &= 2\delta_{2+}\delta_{1+}\frac{\kappa^*}{x_3}\left[\pi_3^* S_+\delta_+ + (1-x_3) m S_-\delta_-\right] \ .
\end{aligned}$$

References

[1] E.A. Kuraev, A. Schiller, V.G. Serbo, D.V. Serebryakova, Eur. Phys. J. C 4 (1998) 631.

[2] E.A. Kuraev, A. Schiller, V.G. Serbo, Nucl. Phys. B 256 (1985) 189; Phys. Lett. B 134 (1984) 455.

[3] E.A. Kuraev, A. Schiller, V.G. Serbo, Z. Phys. C 30 (1986) 237.

[4] A.I. Akhiezer, V.B. Berestetskii, Quantum electrodynamics, Moscow: Nauka 1981.

[5] V.N. Baier, V.M. Galitsky, Phys. Lett. 13 (1964) 355.

[6] E.A. Kuraev et al., Yad. Fiz. 19 (1974) 331.

[7] E.A. Kuraev, V.S. Fadin, Novosibirsk preprint INP 78-56 (1978);
V.N. Baier et al., Phys. Rep. 78 (1981) 293.

[8] E.A. Kuraev, A.N. Peryshkin, Yad. Fiz. 42 (1985) 1195.

[9] F.A. Behrends et al., Nucl. Phys. B 264 (1986) 265.

[10] S. Jadach et al., CERN Yellow Report CERN 95-03 (1995) 343; A.B. Arbuzov et al., ibid. 369; M. Cacciari et al., ibid. 389.

[11] V.B. Berestetskii, E.M. Lifshitz, L.P. Pitaevskii, Quantum electrodynamics, Moscow: Nauka 1989.

[12] E.A. Kuraev, L.N. Lipatov, M.I. Strikman, Zh. Eksp. Teor. Fiz. 66 (1974) 838.